

=> d his

(FILE 'HOME' ENTERED AT 11:26:25 ON 12 SEP 2002)

FILE 'MEDLINE' ENTERED AT 11:27:00 ON 12 SEP 2002

E ZAMBIDIS E/AU

L1 4 S E3-E4
L2 4 DUP REM L1 (0 DUPLICATES REMOVED)

FILE 'STNGUIDE' ENTERED AT 11:28:28 ON 12 SEP 2002

FILE 'MEDLINE' ENTERED AT 11:31:35 ON 12 SEP 2002

E SCOTT E/AU

L3 665 S E3-E41
L4 12 S L3 AND TOLERA?
L5 12 DUP REM L4 (0 DUPLICATES REMOVED)
L6 12 S L5
L7 4 S L2
L8 12 S L5 NOT L2
L9 0 S L1 AND L3
E SCOTT D/AU
L10 2516 S E3-E51
L11 173 S L10 AND TOLERA?
L12 11 S L11 AND FUSION
L13 114 S L11 NOT PY>1994

FILE 'STNGUIDE' ENTERED AT 11:48:16 ON 12 SEP 2002

L14 1 S TOLERANCE OR TOLEROGEN OR TOLEROGENIC

FILE 'MEDLINE, BIOSIS, CAPLUS' ENTERED AT 11:50:36 ON 12 SEP 2002

L15 295085 S TOLERANCE OR TOLEROGEN OR TOLEROGENIC
L16 378100 S FUSION
L17 218 S L15 AND L16 AND (IGG OR IMMUNOGLOBULIN OR FC OR VARIABLE)
L18 199 S L17 AND (CELL OR CELLS OR LYMPHOCYTE OR LYMPHOCYTES OR PBL OR
L19 17 S L18 NOT PY>1994
L20 13 DUP REM L19 (4 DUPLICATES REMOVED)

=>

From: Wilson, Michael
Sent: Thursday, September 12, 2002 12:10 PM
To: STIC-ILL
Subject: art req. 09/160076

TI Genetically transferred central and peripheral immune tolerance via retroviral-mediated expression of immunogenic epitopes in hematopoietic progenitors or peripheral B lymphocytes.

AU Zambidis E T; Kurup A; Scott D W

SO MOLECULAR MEDICINE, (1997 Mar) 3 (3) 212-24.

TI Both resting and activated B lymphocytes expressing engineered peptide-Ig molecules serve as highly efficient tolerogenic vehicles in immunocompetent adult recipients.

AU Zambidis E T; Barth R K; Scott D W

SO JOURNAL OF IMMUNOLOGY, (1997 Mar 1) 158 (5) 2174-82.

TI B cell tolerance induction by cross-linking of membrane IgM, but not IgD, and synergy by cross-linking of both isotypes.

AU Gaur A; Yao X R; Scott D W

SO JOURNAL OF IMMUNOLOGY, (1993 Mar 1) 150 (5) 1663-9.

TI Superantigen modulation of experimental allergic encephalomyelitis: activation of anergy determines outcome.

AU Racke M K; Quigley L; Cannella B; Raine C S; McFarlin D E; Scott D

SO JOURNAL OF IMMUNOLOGY, (1994 Feb 15) 152 (4) 2051-9.

TI Analysis of B cell tolerance in vitro.

AU Scott D W

SO ADVANCES IN IMMUNOLOGY, (1993) 54 393-425. Ref: 141

TI Pathways and regulation of B-cell responsiveness and tolerance.

AU Scott D W; Warner G L; Yao X R; Kent S C

SO CHEMICAL IMMUNOLOGY, (1994) 58 34-66. Ref: 147

TI Long-term survival of xenogeneic pancreatic islet grafts induced by

AU Lenschow D J; Zeng Y; Thistlethwaite J R; Montag A; Brady W; Gibson M G;

SO SCIENCE, (1992 Aug 7) 257 (5071) 789-92.

TI A soluble divalent class I major histocompatibility complex molecule inhibits alloreactive T cells at nanomolar concentrations.

From: Wilson, Michael
Sent: Thursday, September 12, 2002 12:10 PM
To: STIC-ILL
Subject: art req. 09/160076

TI Genetically transferred central and peripheral immune tolerance via retroviral-mediated expression of immunogenic epitopes in hematopoietic progenitors or peripheral B lymphocytes.

AU Zambidis E T; Kurup A; Scott D W

SO MOLECULAR MEDICINE, (1997 Mar) 3 (3) 212-24.

TI Both resting and activated B lymphocytes expressing engineered peptide-Ig molecules serve as highly efficient tolerogenic vehicles in immunocompetent adult recipients.

AU Zambidis E T; Barth R K; Scott D W

SO JOURNAL OF IMMUNOLOGY, (1997 Mar 1) 158 (5) 2174-82.

TI B cell tolerance induction by cross-linking of membrane IgM, but not IgD, and synergy by cross-linking of both isotypes.

AU Gaur A; Yao X R; Scott D W

SO JOURNAL OF IMMUNOLOGY, (1993 Mar 1) 150 (5) 1663-9.

TI Superantigen modulation of experimental allergic encephalomyelitis: activation of anergy determines outcome.

AU Racke M K; Quigley L; Cannella B; Raine C S; McFarlin D E; Scott D

SO JOURNAL OF IMMUNOLOGY, (1994 Feb 15) 152 (4) 2051-9.

TI Analysis of B cell tolerance in vitro.

AU Scott D W

SO ADVANCES IN IMMUNOLOGY, (1993) 54 393-425. Ref: 141

TI Pathways and regulation of B-cell responsiveness and tolerance.

AU Scott D W; Warner G L; Yao X R; Kent S C

SO CHEMICAL IMMUNOLOGY, (1994) 58 34-66. Ref: 147

TI Long-term survival of xenogeneic pancreatic islet grafts induced by

AU Lenschow D J; Zeng Y; Thistlethwaite J R; Montag A; Brady W; Gibson M G;

SO SCIENCE, (1992 Aug 7) 257 (5071) 789-92.

TI A soluble divalent class I major histocompatibility complex molecule inhibits alloreactive T cells at nanomolar concentrations.

STIC-ILL

From:
Sent:
T :
Subject:

Wilson, Michael
Thursday, September 12, 2002 12:10 PM
STIC-ILL
art req. 09/160076

OK out 10/9/12

4/12/01

TI Genetically transferred central and peripheral immune tolerance via retroviral-mediated expression of immunogenic epitopes in hematopoietic progenitors or peripheral B lymphocytes.

AU Zambidis E T; Kurup A; Scott D W

SO MOLECULAR MEDICINE, (1997 Mar) 3 (3) 212-24.

TI Both resting and activated B lymphocytes expressing engineered peptide-Ig molecules serve as highly efficient tolerogenic vehicles in immunocompetent adult recipients.

AU Zambidis E T; Barth R K; Scott D W

SO JOURNAL OF IMMUNOLOGY, (1997 Mar 1) 158 (5) 2174-82.

8101992
Scientific and Technical
Information Center

TI B cell tolerance induction by cross-linking of membrane IgM, but not IgD, and synergy by cross-linking of both isotypes.

AU Gaur A; Yao X R; Scott D W

SO JOURNAL OF IMMUNOLOGY, (1993 Mar 1) 150 (5) 1663-9.

SEP 13 RECD
PAT. & TM OFFICE
COMPLETED

TI Superantigen modulation of experimental allergic encephalomyelitis: activation of anergy determines outcome.

AU Racke M K; Quigley L; Cannella B; Raine C S; McFarlin D E; Scott D

SO JOURNAL OF IMMUNOLOGY, (1994 Feb 15) 152 (4) 2051-9.

TI Analysis of B cell tolerance in vitro.

AU Scott D W

SO ADVANCES IN IMMUNOLOGY, (1993) 54 393-425. Ref: 141

TI Pathways and regulation of B-cell responsiveness and tolerance.

AU Scott D W; Warner G L; Yao X R; Kent S C

SO CHEMICAL IMMUNOLOGY, (1994) 58 34-66. Ref: 147

8379466

TI Long-term survival of xenogeneic pancreatic islet grafts induced by

AU Lenschow D J; Zeng Y; Thistlethwaite J R; Montag A; Brady W; Gibson M G;

SO SCIENCE, (1992 Aug 7) 257 (5071) 789-92.

TI A soluble divalent class I major histocompatibility complex molecule inhibits alloreactive T cells at nanomolar concentrations.

STIC-ILL

From: Wilson, Michael
Sent: Thursday, September 12, 2002 12:10 PM
To: STIC-ILL
Subject: art req. 09/160076

YR 10/9/12

4/21/02

TI Genetically transferred central and peripheral immune tolerance via retroviral-mediated expression of immunogenic epitopes in hematopoietic progenitors or peripheral B lymphocytes.

AU Zambidis E T; Kurup A; Scott D W

SO MOLECULAR MEDICINE, (1997 Mar) 3 (3) 212-24.

TI Both resting and activated B lymphocytes expressing engineered peptide-Ig molecules serve as highly efficient tolerogenic vehicles in immunocompetent adult recipients.

AU Zambidis E T; Barth R K; Scott D W

SO JOURNAL OF IMMUNOLOGY, (1997 Mar 1) 158 (5) 2174-82.

8/6/983

TI B cell tolerance induction by cross-linking of membrane IgM, but not IgD, and synergy by cross-linking of both isotypes.

AU Gaur A; Yao X R; Scott D W

SO JOURNAL OF IMMUNOLOGY, (1993 Mar 1) 150 (5) 1663-9.

TI Superantigen modulation of experimental allergic encephalomyelitis: activation of anergy determines outcome.

AU Racke M K; Quigley L; Cannella B; Raine C S; McFarlin D E; Scott D

SO JOURNAL OF IMMUNOLOGY, (1994 Feb 15) 152 (4) 2051-9.

TI Analysis of B cell tolerance in vitro.

AU Scott D W

SO ADVANCES IN IMMUNOLOGY, (1993) 54 393-425. Ref: 141

TI Pathways and regulation of B-cell responsiveness and tolerance.

AU Scott D W; Warner G L; Yao X R; Kent S C

SO CHEMICAL IMMUNOLOGY, (1994) 58 34-66. Ref: 147

TI Long-term survival of xenogeneic pancreatic islet grafts induced by

AU Lenschow D J; Zeng Y; Thistlethwaite J R; Montag A; Brady W; Gibson M G;

SO SCIENCE, (1992 Aug 7) 257 (5071) 789-92.

TI A soluble divalent class I major histocompatibility complex molecule inhibits alloreactive T cells at nanomolar concentrations.

8011158

Fr m: *AM 1632*
Sent: Wilson, Michael
T: Thursday, September 12, 2002 12:10 PM
Subject: STIC-ILL
art req. 09/160076

TI Genetically transferred central and peripheral immune tolerance via retroviral-mediated expression of immunogenic epitopes in hematopoietic progenitors or peripheral B lymphocytes.

AU Zambidis E T; Kurup A; Scott D W

SO MOLECULAR MEDICINE, (1997 Mar) 3 (3) 212-24.

TI Both resting and activated B lymphocytes expressing engineered peptide-Ig molecules serve as highly efficient tolerogenic vehicles in immunocompetent adult recipients.

AU Zambidis E T; Barth R K; Scott D W

SO JOURNAL OF IMMUNOLOGY, (1997 Mar 1) 158 (5) 2174-82.

TI B cell tolerance induction by cross-linking of membrane IgM, but not IgD, and synergy by cross-linking of both isotypes.

AU Gaur A; Yao X R; Scott D W

SO JOURNAL OF IMMUNOLOGY, (1993 Mar 1) 150 (5) 1663-9.

TI Superantigen modulation of experimental allergic encephalomyelitis: activation of anergy determines outcome.

AU Racke M K; Quigley L; Cannella B; Raine C S; McFarlin D E; Scott D

SO JOURNAL OF IMMUNOLOGY, (1994 Feb 15) 152 (4) 2051-9.

TI Analysis of B cell tolerance in vitro.

AU Scott D W

SO ADVANCES IN IMMUNOLOGY, (1993) 54 393-425. Ref: 141

TI Pathways and regulation of B-cell responsiveness and tolerance.

AU Scott D W; Warner G L; Yao X R; Kent S C

SO CHEMICAL IMMUNOLOGY, (1994) 58 34-66. Ref: 147

TI Long-term survival of xenogeneic pancreatic islet grafts induced by

AU Lenschow D J; Zeng Y; Thistlethwaite J R; Montag A; Brady W; Gibson M G;

SO SCIENCE, (1992 Aug 7) 257 (5071) 789-92.

TI A soluble divalent class I major histocompatibility complex molecule inhibits alloreactive T cells at nanomolar concentrations.

From: Wilson, Michael
Sent: Thursday, September 12, 2002 12:10 PM
To: STIC-ILL
Subject: art req. 09/160076

TI Genetically transferred central and peripheral immune tolerance via retroviral-mediated expression of immunogenic epitopes in hematopoietic progenitors or peripheral B lymphocytes.

AU Zambidis E T; Kurup A; Scott D W

SO MOLECULAR MEDICINE, (1997 Mar) 3 (3) 212-24.

TI Both resting and activated B lymphocytes expressing engineered peptide-Ig molecules serve as highly efficient tolerogenic vehicles in immunocompetent adult recipients.

AU Zambidis E T; Barth R K; Scott D W

SO JOURNAL OF IMMUNOLOGY, (1997 Mar 1) 158 (5) 2174-82.

TI B cell tolerance induction by cross-linking of membrane IgM, but not IgD, and synergy by cross-linking of both isotypes.

AU Gaur A; Yao X R; Scott D W

SO JOURNAL OF IMMUNOLOGY, (1993 Mar 1) 150 (5) 1663-9.

TI Superantigen modulation of experimental allergic encephalomyelitis: activation of anergy determines outcome.

AU Racke M K; Quigley L; Cannella B; Raine C S; McFarlin D E; Scott D

SO JOURNAL OF IMMUNOLOGY, (1994 Feb 15) 152 (4) 2051-9.

TI Analysis of B cell tolerance in vitro.

AU Scott D W

SO ADVANCES IN IMMUNOLOGY, (1993) 54 393-425. Ref: 141

TI Pathways and regulation of B-cell responsiveness and tolerance.

AU Scott D W; Warner G L; Yao X R; Kent S C

SO CHEMICAL IMMUNOLOGY, (1994) 58 34-66. Ref: 147

TI Long-term survival of xenogeneic pancreatic islet grafts induced by

AU Lenschow D J; Zeng Y; Thistlethwaite J R; Montag A; Brady W; Gibson M G;

SO SCIENCE, (1992 Aug 7) 257 (5071) 789-92.

TI A soluble divalent class I major histocompatibility complex molecule inhibits alloreactive T cells at nanomolar concentrations.